

Woven coronary artery: a case report and literature review

S.-M. Yuan

1st Hospital of Putian, Teaching Hospital, Fujian Medical University, Putian, Fujian Province, People's Republic of China

[Received 9 April 2013; Accepted 17 May 2013]

Woven coronary artery is extremely rare. It is characterised by thin channels arising from the coronary artery and reanastomosis at the distal portion. A 62-year-old man was diagnosed of coronary artery disease. Coronary angiography showed 3-vessel coronary artery disease. The distal right coronary artery derived 3 twisting thin channels, and the inferior thin channel sprouted second-class thin channels, which then reanastomosed distally. He received off-pump coronary artery bypass. The present patient had woven coronary artery with a more complex configuration of thin channels different from the previously reported cases. (Folia Morphol 2013; 72, 3: 263–266)

Key words: coronary artery disease, off-pump coronary artery bypass, internal mammary-coronary artery anastomosis

INTRODUCTION

Woven coronary artery is extremely rare. It is characterised by thin channels arising from the coronary artery and reanastomosis at the distal portion of the coronary artery [8]. Woven coronary artery is a benign condition, usually with normal blood flow through the coronary artery [9]. It is interpreted as a congenital anomaly of the coronary artery. However, the incidence of this condition is undetermined [8]. Only 10 patients were reported to have woven coronary artery in the medical literature. In this report, an alternative woven coronary artery was supplemented and the clinical implications of this condition were briefly discussed.

CASE REPORT

A 62-year-old man was referred to our hospital due to chest distress and precordial pain for 5 days. Physical examination revealed that his blood pressure was 180/110 mm Hg on admission. His blood troponin T was 9.26 ng/mL (normal 0–0.04 ng/mL), hypersensitive C-reactive protein was 5.90 mg/L (normal 0–3.0 mg/L), and β_2 -microglobulin was 3.34 mg/L (normal 0.8–3.16 mg/L). Rales were audible

over the lungs, and a 2/6 grade diastolic murmur was heard at the apex. Electrocardiogram showed ST depression in leads II, III, aVF and V4–V6. Echo-cardiography demonstrated left ventricular hypokinesis in the posterior and inferior walls and mild mitral insufficiency with an ejection fraction of 54%. Coronary angiography showed the distal left main coronary artery 50% stenosis, diffuse narrowing of the left anterior descending coronary artery 30 mm long with a maximal 95% stenosis, and proximal right coronary artery 50% diffuse stenosis. The distal portion of the right coronary artery derived three thin channels which were twisting but not intertwining, and the inferior thin channel sprouted second-class thin channels, which then reanastomosed distally (Fig. 1). After medical treatment, the patients' symptoms were relieved, and his blood pressure was normal. He received off-pump coronary artery bypass with a left internal mammary artery-saphenous vein I-graft to the left anterior descending coronary artery, and a sequential saphenous vein grafts to the diagonal and first obtuse marginal branches. He was doing well after the operation.




Figure 1. Coronary angiogram showed (A) long narrowing of left main coronary artery (1), severe stenosis of the left anterior descending coronary artery (2) and of the circumflex branch (3); B, C. Woven right coronary artery derived 3 thin channels with the inferior one sending out second-class thin channels (dashed arrows), which then reanastomosed distally (solid arrow).

DISCUSSION

Ten patients with woven coronary artery were reported in the literature [1–7, 9]. There were totally 11 patients including the present case. Their demographic data were listed in Table 1. This patient setting was comprised of 10 adults and one 9-month-old infant. Their mean age was 49.1 ± 17.4 years (median: 55 years; range: 9 months to 62 years). Eight patients were males, 2 were females and 1 patient's gender was not given.

There were 14 segments of woven coronary arteries in the 11 patients: 10 (90.9%) patients had 1 woven segment, and 1 (9.1%) patient had 4 woven segments. The diagnoses were made on coronary angiography in all 11 patients. The woven coronary arteries were 7 (50%) right coronary arteries, 4 (28.6%) circumflex branches, 2 (14.3%) left anterior descending coronary arteries, and 1 (7.1%) left obtuse marginal branch. The woven segments were located in the distal, proximal, mid, mid-distal, and proximal-mid portion of the coronary arteries in 5 (35.7%), 4 (28.6%), 2 (14.3%), 2 (14.3%) and 1 (7.1%), respectively. The woven coronary arteries were configured as intertwined in 7 (50%), twisting in 5 (35.7%), paralleling in 1 (7.1%), and in a fashion of "figure of 8" in 1 (7.1%), respectively. The thin channels of the woven coronary arteries were double in 8 (57.1%), multiple in 5 (35.7%), and triple in 1 (7.1%) segment of the coronary arteries, respectively. The woven coronary arteries were with normal flow in all 11 patients. Occlusion of otherwise coronary arteries developed in 8 patients and coronary artery dilation secondary to Kawasaki disease in

1 patient. Four of the patients with coronary artery occlusion received percutaneous coronary angioplasty and 2 patients underwent coronary artery bypass grafting. All patients survived at a maximal follow-up of 4 years.

Woven coronary arteries often have thin channels and reanastomosis at the distal part of the coronary artery. However, in the exceptional case, reanastomosis may be absent at the farther distal portion of the coronary artery [2]. Coronary angiography is the gold standard for the diagnosis of woven coronary artery [8]. In spite of abnormal coronary artery structures, the blood flow and the myocardial contractility are normal [5, 8]. The thin channels were occasionally misdiagnosed as a dissected atherosclerotic plaque with thrombus formation [4, 6]. Kaya et al. [4] reported that the thin channels were not visualised until the third coronary angiography 3 years after coronary artery bypass grafting.

The present patient had woven coronary artery with a more complex configuration of thin channels different from the previously reported cases.

CONCLUSIONS

Woven coronary artery is an extremely rare lesion. The thin channels that the woven coronary arteries form may sometimes lead to a misdiagnosis. As woven coronary artery is not a substantially stenosed coronary artery and therefore does not require special treatment, care must be taken when the physicians encounter such a lesion and differential diagnosis with a dissected atherosclerotic plaque with thrombus formation is warranted.

Table 1. Clinical features of the patients with woven coronary artery

Year	Author	Age	Gender	Symptom	ECG	ECHO	Thallium scintigraphy	Woven coronary artery		Diseased coronary artery	Management	Outcome
								Location	Thin channels			
								No.	Configuration	Length [cm]		
1988	Sane and Vidaillet	55	F	Congestive heart failure				Proximal RCA	2	Figure of 8		
1990	Berman et al.	51	M	No	ST depressions			Mid-distal RCA	2	Intertwined	PDA	
1995	Gregorini et al.	62	M	Angina, s/p AMI				Proximal-mid Cx	2	Intertwined	LAD, Cx	PTCA on mid-LAD
1995	Gregorini et al.	60	?	Angina pectoris			Positive	LAD, Cx, OM	3 in LAD, 2 in Cx and OM	Intertwined		
1995	Gregorini et al.	45	F	Angina, AMI				Distal LAD	2	Parallel	LAD	
2000	Martuscelli et al.	42	M	Angina				Mid-RCA	2	Intertwined	LAD, OM ₁	LAD stenting, OM ₁ stenting
2006	Kursaklioglu et al.	48	M	Exertional chest pain	Normal (resting), ST segment depression (stress)	Moderate aortic insufficiency		Mid-Cx	Multiple	Twisting	RCA	RCA stenting, AVR
2006	Kaya et al.	56	M	Stable angina pectoris, exercise dyspnoea				Mid-distal RCA	Multiple	Slight twisting	LM, LAD	CABG × 2: LIMA→LAD, SVG→OM ₁
2010	Yildirim et al.	0.75	M	Kawasaki disease, high fever	Marked PR interval			Proximal RCA	Multiple	Slight twisting	LAD dilation	Conservative treatment
2010	Iyisoy et al.	58	M	Exertional chest pain	ST depression in leads V4–V6 (stress)	Normal left ventricular wall motion	Reversible ischemia in the anterior wall	Proximal RCA	Multiple	Slight twisting	1 LAD	LAD stenting
2013	Yuan	62	M	Angina pectoris, s/p AMI	ST depression in leads II, III, aVF and V4–V6	LV hypokinesis, mild mitral insufficiency		Distal RCA	Multiple	Twisting, not intertwined	LM, LAD, Cx, RCA	LAD 70% stenosis 3 years after stenting

AMI — acute myocardial infarction; AVR — aortic valve replacement; CABG — coronary artery bypass grafting; Cx — circumflex branch of the coronary artery; ECG — electrocardiogram; ECHO — echocardiography; F — female; LAD — left anterior descending coronary artery; LIMA — left internal mammary artery; LM — left main coronary artery; LV — left ventricle; M — male; MVC — mitral valve commissurotomy; OM — obtuse marginal branch; PDA — posterior descending coronary artery; PTCA — percutaneous transluminal coronary angioplasty; RCA — right coronary artery; s/p — status post; SVG — saphenous vein graft

REFERENCES

1. Berman AD, Kim D, Baim DS (1990) "Woven" right coronary artery: case report and therapeutic implications. *Cathet Cardiovasc Diagn*, 21: 258–259.
2. Gregorini L, Perondi R, Pomidossi G, Saino A, Bossi IM, Zanchetti A (1995) Woven left coronary artery disease. *Am J Cardiol*, 75: 311–312.
3. Iyisoy A, Celik T, Yuksel UC, Isik E (2010) Woven right coronary artery: a case report and review of the literature. *Clin Cardiol*, 33: E43–E45.
4. Kaya D, Kilit C, Onrat E (2006) An uncommon congenital anomaly of coronary arteries misdiagnosed as intracoronary thrombus: woven coronary artery disease. *Anadolu Kardiyol Derg*, 6: 383–384.
5. Kursaklioglu H, Iyisoy A, Celik T (2006) Woven coronary artery: a case report and review of literature. *Int J Cardiol*, 113: 121–123.
6. Martuscelli E, Romeo F, Giovannini M, Nigri A (2000) Woven coronary artery: differential diagnosis with diffuse intracoronary thrombosis. *Ital Heart J*, 1: 306–307.
7. Sane DC, Vidaillet HJ Jr. (1988) "Woven" right coronary artery: a previously undescribed congenital anomaly. *Am J Cardiol*, 61: 1158.
8. Yıldırım A, Oğuz D (2011) Woven coronary artery. In: Suna F. Kiraç ed. *Advances in the diagnosis of coronary atherosclerosis*. InTech, 297–304. Available from: <http://www.intechopen.com/articles/show/title/woven-coronary-artery>.
9. Yıldırım A, Oğuz D, Olguntürk R (2010) Woven right and aneurysmatic left coronary artery associated with Kawasaki disease in a 9-month-old patient. *Cardiol Young*, 20: 342–344.